



Do Neurological Changes in CT Brain in Eclamptic Patients Correlate With Neurological Symptoms? : An Observational Study

¹T.H.Usha*, ²S.Valarmathi,

ABSTRACT

To evaluate the different neurological changes in brain in eclampsia by CT scan in relation to neurologic symptoms. This is prospective observational study in our hospital. CT scan of brain is performed within 48 hours of eclampsia after confinement of fetus and after stabilising the mother with standard MgSO₄ protocol. The CT scans of brain are performed with 5mm and 10mm section in the axial plain. CT scan of brain shows, 32% have cerebral edema, 23% have cerebral infarct, 8% have cerebral haemorrhage, while 37% have no detectable findings. Parietal region of the brain is affected in 67% followed by parieto-occipital area (17%), occipital area (8%) and brain stem (8%). 68 % mothers have headache, 18% have visual disturbances, 34% have altered sensorium with hyper-reflexia and 37% have coma. CT scan of brain in eclampsia can provide useful intra cerebral information and should be done in cases with severe neurologic manifestations, if possible for every eclamptic mother.

KEY WORDS : Eclampsia, Cerebral Edema, Cerebral Hemorrhage, Cerebral Infarct, Altered Sensorium, Visual Disturbance, Coma.

Introduction

Eclampsia is defined as occurrence of generalised seizures, not caused by any co-incidental neurological disorder (e.g. epilepsy) in a woman whose condition also meets the criteria for preeclampsia [1] which is a complex multi-organ disorder characterised by pregnancy induced hypertension and proteinuria after 20 wks of pregnancy (exception—gestational trophoblastic disease or multiple pregnancy).

Cerebral complications are the major cause of deaths in eclampsia; still the neuro pathophysiology of eclamptic seizure is mostly unknown. There are two distinct but related types of cerebral pathology in the patients of eclampsia [2]. The first is gross haemorrhage due to ruptured arteries caused by severe hypertension of any cause, not necessarily only by preeclampsia or eclampsia. The second type of post-mortem lesions are edema, hyperaemia, ischemic micro infarcts and petechial haemorrhages. The neurologic manifestations of severe eclampsia are identical to those of hypertensive encephalopathy [2], which is clinically manifested as generalised tonic-clonic seizure and usually preceded by neurological symptoms like hyper-reflexia, altered sensorium, headache, visual changes and even coma.

^{1,2}Senior Assistant Professor of Obstetrics and Gynaecology, Government Rajaji Hospital, Madurai Medical College, Madurai.

*Corresponding Author

Dr. T.H.Usha,
Senior Assistant Professor of Obstetrics and Gynaecology,
Government Rajaji Hospital,
Madurai Medical College,
Madurai-625020,
Mail id: ushadrbalan@gmail.com
Mobile.No: +919443587008

The recent advances in radiologic imaging including the use of computed tomography (CT) scans and magnetic resonance imaging (MRI), have greatly enhanced our understanding about the correlation between neurologic manifestations and neuro-anatomic and pathological characteristics of eclampsia [3]. Harandou M et al [4]; showed that 73.68% cases of eclamptic mothers who are still symptomatic after 24 hours have cerebral edema and 10.5% have cerebral haemorrhage and 15.7% have normal CT scan study.

The aim of the study is to evaluate the different neurological changes in brain by CT scan in eclampsia and their relation with different neurologic symptoms. In this study, CT scan methodology has been adopted because it is less expensive and easily available.

Methodology

This is a prospective study of CT scan finding of brain on cases of eclampsia admitted in our hospital. The study population are chosen by random samplings who are patient of eclampsia admitted through emergency and also indoor patients who develop eclampsia after admission.

Inclusion Criteria

Patients with Eclampsia (at least one episode of seizure in women with more than 20 weeks gestation or less than 06 weeks postpartum with blood pressure more than 140 mm of Hg systolic and 90 mm of Hg diastolic with urine albumin of more than 0.3gm/L). Both ante partum and postpartum.

Exclusion Criteria

1. Women who are known case of Hypertension, Epilepsy.
2. Seizures due to metabolic disturbances, space occupying lesions or intra cerebral infections.

Total 100 eclamptic mothers are chosen according to inclusion criteria. Study period is of one year and 9 months from Jan 2014 to Sep 2015. Basic information including age, parity and gestational age, previous medical or obstetric history is taken. Detailed history of convulsion like duration, time, number of convulsion and presence of premonitory symptoms are sought; followed by detailed neurological examination (specially level of consciousness, pupillary reaction and reflexes) including fundoscopy is performed. Basic investigations like blood pressure, urine for proteinuria (by dipstick) are measured and complete hemogram, platelet count, serum uric acid, serum creatinine, liver enzymes are sent. Standard MgSO₄ protocol is given to all eclamptic mothers.

If the mother is not already delivered, assessment of cervix and delivery of the fetus is done accordingly either by induction of labour or Caesarean section. CT scan of brain is performed within 48 hours of eclampsia after confinement of fetus and after stabilising the mother. The CT scans of brain are performed with plain and intravenous (non-ionic) contrast enhancement (if necessary) with 5mm and 10mm section in the axial plain. The CT scan findings are evaluated with neurological characteristics. Level of consciousness is classified according to Glasgow coma scale (<8 severe, 9-12 moderate and >13 minor) [5]. If CT scan of brain is normal but if has persistent neurological symptoms MRI BRAIN with MRA and MRV is suggested. MRI is not done in all as it is not easily available and it being expensive. Statistical analysis is performed with aid of Statistical Package for the Social Sciences. P value <0.05 is considered for statistical significance. Follow-up CT scan is not performed as it is not included in the study protocol.

Results

Total 100 eclamptic mothers are included in this study. Median age of the mothers is 23 years with standard deviation (SD) of 3.8 years. 55% of eclamptic mothers are primigravida and 45% of eclamptic mothers are multigravida. Among them 28% have postpartum eclampsia, 4% have intra partum eclampsia and 68% have ante partum eclampsia. 35% mothers delivered by normal delivery and 65% mothers have undergone LSCS.

CT scan of brain shows, 32% have cerebral edema (diffuse white matter low density areas, patchy area of low density, loss of normal cortical sulci) 23% have cerebral infarct (hypo attenuating brain tissue), 8% have cerebral haemorrhage (intra ventricular/ parenchymal haemorrhage, subarachnoid hemorrhage, subdural hematoma), while 37% have no detectable findings. (Table.No.1) Parietal region of the brain is affected in 67% followed by parieto-occipital area (17%), occipital area (8%) and brain stem (8%) (Table.No. 2).

Among different neurologic symptoms 68% mothers have headache, 18% have visual disturbances, 34% have altered sensorium with hyper-reflexia and 37% have coma. Eclamptic mother who presented with visual disturbances (18/100) mostly have brain lesions in parieto-occipital and occipital region (16/18), which is statistically significant ($p < 0.005$). Similarly, mothers presented with coma (37/100) mostly have lesions in parietal cortex (26/37) also, significant ($p = 0.002$). But no association is found with area of lesions and other symptoms like headache. Hyperreflexia indicates pyramidal syndrome involving CNS but has no correlation with type of lesions. 53% eclamptic mothers are preterm (<37 weeks completed gestational age); among them 43% have cerebral edema, 29% have cerebral infarction, 14% have cerebral haemorrhage and 14% have no CT scan findings. 47% eclamptic mothers are

term (>37 weeks completed gestational age); among them 18% have cerebral edema, 18% have cerebral infarction, but 64% have no CT scan findings.

In this study there is no difference between blood pressure distributions between those who have CT scan findings than those who have not positive CT scan findings. (Table. No.3)

Eclamptic mother whose number of episode of convulsion is less than 5; among them 63% have no finding in CT scan, 28% cerebral edema, 9% have cerebral infarction. On the other hand whose number of episode of convulsion is more than 5, among them 39% develop cerebral edema, 44% develop infarction and 17% develop cerebral haemorrhage ($P = 0.001$). 55% eclamptic mothers have Glasgow coma scale <8 during admission; among them 33% develop cerebral edema, 33% develop infarction, 15% develop cerebral haemorrhage and 19% have no CT scan findings. 45% eclamptic mother whose Glasgow coma scale is >8; among them 29% develop cerebral edema, 12% develop infarct and 59% have no CT scan finding ($p < 0.05$). (Table. No. 4) Of the eclamptic mothers who recovered within 24 hours to fully oriented state ($N=37$) 77% have no CT scan findings, only 16% develop cerebral edema and 7% develop infarction. Eclamptic mother who recovered over 48 hours ($N=37$); only 10% have no CT scan finding in CT scan, 46% develop edema, 32% develop infarction and 12% develop hemorrhage ($p < 0.019$).

Table. No. 1: CT scan findings in eclamptic patients

| CT finding | Percentage of patients (n=100) |
|-----------------------|--------------------------------|
| Cerebral edema | 32 |
| Cerebral infarct | 23 |
| Cerebral hemorrhage | 08 |
| No detectable finding | 37 |
| Total | 100 |

Table.No. 2: Different areas of brain involvement by CT brain

| | Cerebral odema | Cerebral hemorrhage | Cerebral infarct | Total |
|------------------------------------|----------------|---------------------|------------------|---------|
| Basal Ganglia and internal capsule | 2 | 0 | 3 | 5(8%) |
| Occipital cortex | 5 | 0 | 0 | 5(8%) |
| Parietal cortex | 22 | 5 | 15 | 42(67%) |
| Parieto-occipital cortex | 3 | 3 | 5 | 11(17%) |

Table.No. 3: BP distribution among eclamptic mothers

| | CT scan features (Edema / Hemorrhage/Infarction) | No CT scan findings | P value |
|------------------------|--|---------------------|---------|
| Systolic BP (Mean±SD) | 166.25+17.64mmHg | 155.71+17.85mmHg | 0.086 |
| Diastolic BP (Mean±SD) | 113.33+14.09mmHg | 107.14+9.94mmHg | 0.156 |

Table.No. 4: CT Scan findings in different neurologic symptoms

| | CT scan of brain | | | | |
|-----------------------------------|-------------------|----------------|---------------------|---------------------|--|
| | Normal CT finding | Cerebral odema | Cerebral infarction | Cerebral hemorrhage | |
| Neurologic symptoms | | | | | |
| Altered sensorium & hyperreflexia | 38% | 47% | 15% | 0 | |
| Head ache | 26% | 31% | 31% | 12% | |
| Visual disturbance | 14% | 28% | 43% | 15% | |
| Coma | 7% | 35% | 35% | 23% | |

Table.No. 5 CT scan findings in eclamptic patients in different studies.

| | Normal finding | Edema | Infarction /thrombosis | Haemorrhage |
|--------------------|----------------|--------|------------------------|-------------|
| Harandou et al [4] | 15.78% | 73.68% | 15.78% | 10.53% |
| Akan H et al [20] | 18.18% | 50% | 13.63% | 9.09% |
| Milliez J | 59% | 34% | | 6.8% |
| Richards AM[15] | | 63.79% | | 9.3% |
| Naidu K[6] | | 58.5% | | |

Discussion

In this study cerebral edema is most common lesion (32%) detected by CT scan, but most importantly 37% eclamptic mothers have no CT scan finding. These finding is corroborative with the findings of Harandou M et al. [4] and Akan H et al. [6] (Figure.No. 4).

In patients with a normal CT scan, MRI is indicated but has not been done in all even though its given in low cost in our institute. 50% of the Posterior reversible encephalopathy syndrome (PRES) patients (revealed by MRI) show normal initial CT scans. MRI investigation would have revealed more brain lesions.

Regarding area of distribution parietal and occipital area is the most frequent site of brain lesions in CT scan; supported by observation of Naidu et al. [7]. They found parieto-occipital involvement in 97.4% of cases. Sometimes diffuse brain edema is associated with compression or dilatation of 3rd and 4th ventricles. There are two such cases in our study. Two rare cases of lacunar infarct and another one rare subarachnoid haemorrhage is found in this study.

PRES is characterized by headache, altered mental status, visual disturbances, and seizures. Although hypertensive encephalopathy can arise in patients with conditions in which there is acute systemic hypertension alone, it most commonly occurs in patients also having pre-existing endothelial dysfunction or damage. The combination of acute hypertension and endothelial damage results in hydrostatic edema—a specific form of vasogenic edema characterised by the forced leakage of serum through capillary walls and into the brain interstitium— which, if severe enough, will be radio-graphically evident [9-10]. Vasogenic edema is most common finding in eclampsia which explain the reversible nature of most eclampsia. The patients which show no significant finding in CT scan may have very mild vasogenic edema not enough for radiologic detection. The CT scan findings of cerebral infarction are originating from anoxia and cytotoxic edema. This may represents the spectrum of eclampsia ranges from an initially reversible phase of vasogenic edema formation to a later phase of ischemic damage and haemorrhage, which carries a worse prognosis with residual neurologic effect [11]. In fact, laboratory studies of hypertensive encephalopathy, suggest that as vasogenic edema progresses, local tissue pressure increases. This causes a decrease in regional perfusion pressure and a reduction of blood flow to ischemic levels. Subsequently, areas surrounding marked vasogenic edema may progress to infarction and cytotoxic edema [11].

Brain perfusion is maintained by an auto regulatory system of small arteries and arterioles that has myogenic and neurogenic component [10]. In PRES cases direct toxic effect on endothelium or vessel distension decrease the effect of myogenic mechanism. Then neurogenic mechanisms take over regulation of cerebral perfusion. The peri

vascular sympathetic nerves travel in the adventitial layer of cerebral blood vessels and are relatively protected from agents that cause endothelial damage. Since the vertebro-basilar system and posterior cerebral arteries are sparsely innervated by sympathetic nerves [12]; the occipital lobe and other posterior brain regions may be particularly susceptible to breakthrough of auto-regulation with elevated systemic pressure. Vasoconstriction induced by sympathetic innervations, moderately protects anterior circulation areas from over perfusion.

Headache is most common neurologic symptoms in this study (68%). Akutsu T et al (1992) [13] and Chang WN et al (1996) [14] also get similar results. Eclamptic mothers with visual symptoms and coma have more lesions in parieto-occipital region and parietal region respectively is corroborative with the findings of Chakravarty A, Chakrabarty SD (2002) [15] and Chang WN et al (1996) [14]. Mothers who have develop coma with Glasgow coma scale <8 and with recurrent episode of convulsion (>5 times in number) develop more findings in CT scan. This finding is correlated to study of Richards et al [16] showing severity of edema is related to duration of intermittent seizures. Also, mothers who become fully oriented within 24 hours have significantly less chance of having brain lesions in CT scan. As cerebral mass effect along with diffuse white matter hypo-densities is associated significantly more with coma ($p < 0.034$); these mothers recovered later from their eclamptic episodes [17]. In this study preterm eclamptic mother are significantly having pronounced CT scan finding than term mother ($p < 0.05$); as preterm mothers are more severely affected in respect to more prodromal symptoms, multiple seizures, major maternal complication [18]. In our study, there is no statistical significant difference in blood pressure values between cases of positive CT scan findings and cases with normal CT scan findings. Brain edema

detected in preeclampsia/eclampsia is thought to be secondary to endothelial injury, rather than hypertension. This finding is correlated with the findings of Schwartz et al [19].

Conclusion

It is evident from this study that cerebral edema is most common cerebral lesions followed by infarction and haemorrhage and parieto-occipital regions of brain is the most common affected area. Although almost 37% eclamptic mothers do not have cerebral lesions, those who have lesions are significantly related to level of consciousness, number of convulsive episode and time taken to recover fully oriented state. Most common neurological finding is headache followed by altered sensorium and hyper-reflexia, visual disturbances and coma.

CT scan of brain can provide useful intra cerebral information to detect different brain lesions in eclampsia which may have different prognosis with residual effect and may need specific modification in management protocol to prevent long term neurologic sequels and reduce maternal mortality and morbidity; although these parameters are not included in this study.

References

1. Cunningham FG, MacDonald PC, Gant NF. Hypertensive disorder of pregnancy. In: Cunningham FG, MacDonald PC, Gant NF, editors. *Williams' obstetrics*. 18th edn. Norwalk, Conn: Appleton & Lange. 1989; 653-694.
2. Barton JR, Sibai BM. Cerebral pathology in eclampsia. *Clin Perinatol*. 1991; 18: 891-910.
3. Marques R, Braga J, Leite I, Jorge CS. Neurological involvement in pre-eclampsia/eclampsia: the role of neuroimaging. *Acta Med Port*. 1997; 10: 585-588.
4. Harandou M, Madani N, Labibe S, Messouak O, Boujraf S, Benkirane S. Neuroimaging findings in eclamptic patients still symptomatic after 24 hours: a descriptive study about 19 cases]. *Ann Fr Anesth Reanim*. 2006; 25: 577- 583.
5. Teasdale G, Jennett B. Assessment of coma and impaired consciousness. A practical scale. *Lancet*. 1974; 2: 81-84.
6. Akan H, Küçük M, Bolat O, Selçuk MB, Tunali G. The diagnostic value of cranial computed tomography in complicated eclampsia. *J Belge Radiol*. 1993; 76: 304-306.
7. Naidu K, Moodley J, Corr P, Hoffmann M. Single photon emission and cerebral computerised tomographic scan and transcranial Doppler sonographic findings in eclampsia. *Br J Obstet Gynaecol*. 1997; 104: 1165-1172.
8. Schwartz RB, Jones KM, Kalina P, Bajakian RL, Mantello MT, Garada B. Hypertensive encephalopathy: findings on CT, MR imaging, and SPECT imaging in 14 cases. *AJR Am J Roentgenol*. 1992; 159: 379-383.
9. Covarrubias DJ, Luetmer PH, Campeau NG. Posterior reversible encephalopathy syndrome: prognostic utility of quantitative diffusion-weighted MR images. *Am J Neuroradiol*. 2002; 23: 1038-1048.
10. Schwartz RB, Feske SK, Polak JF, DeGirolami U, Iaia A, Beckner KM. Preeclampsia-eclampsia: clinical and neuroradiographic correlates and insights into the pathogenesis of hypertensive encephalopathy. *Radiology*. 2000; 217: 371-376.
11. Koch S, Rabinstein A, Falcone S, Forteza A. Diffusion-weighted imaging shows cytotoxic and vasogenic edema in eclampsia. *Am J Neuroradiol*. 2001; 22: 1068-1070.
12. Beausang-Linder M, Bill A. Cerebral circulation in acute arterial hypertension- protective effects of sympathetic nervous activity. *Acta Physiol Scand*. 1981; 111: 193-199.
13. Akutsu T, Sakai F, Hata T, Iizuka T, Tazaki Y. Neurological and neuroimaging studies of eclampsia. *Rinsho Shinkeigaku*. 1992; 32: 701-707.
14. Chang WN, Lui CC, Chang JM. CT and MRI findings of eclampsia and their correlation with neurologic symptoms. *Zhonghua Yi Xue Za Zhi (Taipei)*. 1996; 57: 191-197.
15. Chakravarty A, Chakrabarti SD. The neurology of eclampsia : some observations. *Neurol India*. 2002; 50: 128-135.
16. Richards A, Graham D, Bullock R. Clinicopathological study of neurological complications due to hypertensive disorders of pregnancy. *J Neurol Neurosurg Psychiatry*. 1988; 51: 416-421.

17. Naheedy MH, Biller J, Schiffer M, Azar-Kia B, Gianopoulous J, Zarandy S. Toxemia of pregnancy: cerebral CT findings. *J Comput Assist Tomogr.* 1985; 9: 497-501.
18. Douglas KA, Redman CW. Eclampsia in the United Kingdom. *BMJ.* 1994; 309: 1395-1400.
19. Schwartz RB, Feske SK, Polak JF, DeGirolami U, Iaia A, Beckner KM. Preeclampsia-eclampsia: clinical and neuroradiographic correlates and insights into the pathogenesis of hypertensive encephalopathy. *Radiology.* 2000; 217: 371-376.
20. Hira B, Moodley J. Role of cerebral computerised tomography scans in eclampsia. *J Obstet Gynaecol.* 2004; 24: 778-779.